## Measurement of Reactive Nitrogen in the Stratosphere.

- G C Toon, 1 3 Sen, G B Osterman, R J Salawitch, J -F Blavier, J J Margitan, 1 D B Peterson IIM 1 lickett, R A Stachnik (Jet 1 ropulsion Laboratory, MS 183-:301, Californial Institute of Technology, 4800" Oak Grove 1 Drive, Pasadena, CA 91 J 09; 818-354-8259; e-mail: toon@mark4sun.jpl.nasa.gov)
- G 4 Yue (NASA Langley Research Center, MS 475, Hampton, VA 23681)

Simultaneous vertical profiles of N(), NO<sub>2</sub>, HNO<sub>3</sub>, HNO<sub>4</sub>, N<sub>2</sub>O<sub>5</sub>, and (HNO<sub>3</sub> were incasured at 35° N over the 20 - 39 km altitude range during a September 1993 balloon flight of the .11'1, MkIV interferometer. We will illustrate the excellent agreement of the measured diurnal variations of NO, NO<sub>2</sub>, and  $2 \times N_2O_5$  with predictions of a photochemical model, constrained by MkIV measurements of NO<sub>y</sub>, Cl<sub>y</sub>, and O<sub>3</sub>. Our results provides confidence that the photochemical model correctly reproduces: (1) the conversion of NO to NO<sub>2</sub> at sunset and (2) the decrease of NO<sub>x</sub> ( $\epsilon$  NO+NO<sub>2</sub>) during nighttime through production of N<sub>2</sub>O<sub>5</sub>.

Simultaneous observations of the suite of NO<sub>y</sub> gases by Mk IV during a series of balloon flights between 1990 and 1993, combined with aerosol surface area from SAGE II, allow us to probe the role of heterogeneous reactions of I sulphate aerosols. The inclusion in the photochemical model of het erogeneous hydrolysis of N<sub>2</sub>O<sub>5</sub> is sufficient to provide an excellent description of the observed non-linear dependence of NO<sub>x</sub>/N O<sub>y</sub> ratio on sulphate aerosol Sulfilite area. Since the saturation of NO<sub>x</sub>/NO<sub>y</sub> ratio limits the increase in ClO within the Cl<sub>y</sub> reservoir, catastrophic loss of O<sub>3</sub> from enhanced sulphate aerosols is precluded for sillifile areas 11) to 14  $\mu$ m<sup>2</sup>.cm<sup>-3</sup>. We will also show that heterogeneous hydrolysis of ClNO<sub>3</sub>, BrNO<sub>3</sub>, and reactions of HCl 011 sulphate aerosols play a minimal role in regulating NO<sub>x</sub> for the observed mid-latitude stratosphere.

- 1. 1996 Fall Meeting
- 2. 010143674
- 3. (a) G C Toon
  Jet Propulsion Laboratory
  M. s. 183-301
  California Inst. of Tech.
  4800 Oak Grove Drive
  Pasadena, CA 91109-8099
  - (b) Tel: (818)354-8259
  - (b) Fax: (818)354-5148
- 4. A
- 5. **(a)** None (1)) 0340"
  - Middle atmosphere composition and chemistry
- 6. Oral Preferred
- 7.
- 8. 100% GRL (submitted)
- 9. Cheque Enclosed
- 10. C
- 11.
- 12.
- 13.